

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 March 2003 (06.03.2003)

PCT

(10) International Publication Number
WO 03/018790 A1

(51) International Patent Classification⁷: C12N 9/24, 15/56, 1/20

(74) Agent: LEE, Hoo, Dong; 7th-11th Floors, Hankook Tire Bldg, 647-15, Yoksam-dong, Gangnam-gu, Seoul 135-723 (KR).

(21) International Application Number: PCT/KR01/01439

(22) International Filing Date: 25 August 2001 (25.08.2001)

(25) Filing Language: Korean

(26) Publication Language: English

(71) Applicant (for all designated States except US): LIFENZA CO., LTD. [KR/KR]; 2nd Floor Samwoo Bldg, 732-24, Yeoksam-dong, Gangnam-gu, Seoul 135-514 (KR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (for US only): KIM, Doman [KR/KR]; Department of Biochemical Engineering, Chonnam National University, 300 Yongbong-dong, Buk-gu, Kwangju-shi 500-757 (KR). KIM, Sung Hyuk [KR/KR]; #506 Woosung Livingtel, 467-10 Dogok-2-dong, Gangnam-gu, Seoul 135-270 (KR). KANG, Hee Kyoung [KR/KR]; Laboratory of Carbohydrate Enzyme and Microbial Engineering, Department of Biochemical Engineering, Chonnam National University, 300 Yongbong-dong, Buk-gu, Kwangju-shi 500-757 (KR).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ENZYME WITH THE REMOVAL



(57) Abstract: The present invention relates to a enzyme of repressing plaque formation and degrading plaque, a DNA sequence encoding the enzyme, the expressing host cell, methods for producing said enzyme, and the isolated and purified enzyme. The invention also relates to compositions comprising the enzyme.

WO 03/018790 A1

ENZYME WITH THE REMOVAL

5

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to an enzyme which can effectively repress formation of plaque or degrade the plaque, gene for encoding the enzyme, and a
10 process for preparing the enzyme and the expressed cell. More specifically, the invention relates to an enzyme which can effectively repress formation of plaque produced from *Lipomyces starkeyi* KSM 22 mutant and degrade the plaque, gene for encoding the enzyme, a process for producing said enzyme and the expressed cells, and a composition comprising the enzyme.

15

Description of the Prior Art

Generally, film formed on surface of a tooth (plaque) consists of intimately packed bacteria and non-cellular substances. The main polysaccharide component of plaque is water-insoluble glucan or mutan, which occupies about
20 20% of total dry weight of plaque, is one of the essential cause of tooth-decay. By way of investigations of the structure of glucan produced by *Streptococcus mutans*, it has been reported that insoluble glucan essentially comprises α -1,3-, α -1,4- and α -1,6-D-glucoside bonds. Thus, the activity of degrading mutan and that of decomposing starch and dextran are required in order to effectively remove
25 plaque.

Conventionally, repression of growth of *Streptococcus mutans* (hereinafter, abbreviated to as "*S. mutans*") has been suggested as a means to repressing plaque

formation or tooth-decay, so that an antibiotic substance to repress growth of *S. mutans* or a fluorine compound was contained in a product for oral use such as mouth washer. Fluorine, as a representative compound of anti-cavity substance, has severe adverse effect of white spot on tooth enamel even at a very low concentration as well as intense toxicity and air pollution, even though it has an effect of repressing growth of bacterial source of tooth-decay. Enzyme such as dextranase has been also used for repressing tooth-decay, but the activity has not been clearly demonstrated yet.

U.S. Pat. 5,741,773 provided a toothpaste composition comprising glycomacropptide having anti-plaque and activity preventing tooth-decay. Such an art, however, simply relates to repression of growth of the bacterial source of tooth-decay, but could not disclose repression of plaque formation or degradation of already formed plaque.

Investigations to employ an enzyme DEXAMase which decomposes polysaccharides of various structures in degradation of already formed plaque have been made by the present inventors. Patent applications regarding a microorganism (*Lipomyces starkeyi* KFCC-11077) producing an enzyme for degrading various polysaccharides, the enzyme and a composition comprising the same are pending in Korea and in the United States.

SUMMARY OF THE INVENTION

The present invention is contrived by said requirements to solve the problems described above.

Accordingly, the present invention has an object to provide novel enzyme having activity to repress formation of plaque or to degrade plaque, and gene encoding the enzyme.

Another object of the present invention is to provide a cell expressing said

gene.

Still another object of the invention is to provide a process for preparing said enzyme and gene.

Still further object of the invention is to provide a commercially useful
5 composition comprised of said enzyme.

In order to achieve the objects, the present invention provide protein comprising amino acid residue of sequence information 1 or mutants thereof, and enzyme that degrades insoluble glucan having both dextranase activity and amylase activity as a part thereof.

10 Further, the present invention provides said protein or its mutants, gene or mutants of sequence information 2 encoding a part thereof, and/or the segments. Preferably, said mutants of protein comprises not less than 80% of homology with amino acid sequence of sequence information 1, and said gene mutants comprises not less than 80% of homology with gene base sequence of sequence information 2.

15 The present invention also provides transformed cells which express said gene. The cells are preferably prokaryotic cell or eukaryotic cell. The most preferable one is E. coli KCTC 10024BP.

The present invention further provides a process preparing a purified enzyme which has both dextranase activity and amylase activity and degrades
20 insoluble glucan, comprising the steps of culturing said cell, expressing an enzyme from the cultured cell, and purifying the expressed enzyme; an enzyme prepared by said process; and a composition comprised of the enzyme.

BRIEF DESCRIPTION OF THE DRAWINGS

25 The present invention will be explained in terms of exemplary embodiments described in detail with reference to the accompanying drawings, which are given only by way of illustration and thus are not limitative of the

present invention:

Fig. 1 shows the effect of pH on activity and stability of DEXAMase: (a) shows the effect on the activity, while (b) shows the effect on the stability.

Fig. 2 shows the effect of temperature on the stability of DEXAMase.

5 Fig. 3 shows a TLC photograph of polysaccharides hydrolyzed by using DEXAMase prepared from recombinant clone pBDEXAM. In the photograph, Mn and IMn shows a series of maltodextrin, lane 1 shows dructose, lane 2 shows dextran, lane 3 shows starch, lane 4 shows inulin, lane 5 shows levan, lane 6 shows mutan, lane 7 shows isomaltose, lane 8 shows maltose, lane 9 shows cellular
10 supernatant, lane 10 shows the TLC photograph after reaction with dextran, lane 11 shows one after reaction with starch, lane 12 shows one after reaction with inulin, lane 13 shows one after reaction with levan, lane 14 shows one after reaction with mutan, lane 15 shows one is after reaction with isomaltose, and lane 16 shows one after reaction with maltose.

15 Fig. 4 shows bonding of DEXAMase having the activity of dextranase and amylase on hydroxyapatite.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Now, the present invention is described in more detail.

20 In order to obtain DEXAMase gene, *L. starkeyi* KSM 22 was cultured in a starch-containing medium, poly(A) + RNA were obtained, cDNA were prepared by the poly(A) + RNA and the cDNA was ligated to Uni-ZAP XR vector. The phage lysate was transformed to pBluescript phagemid via segmenting *in vivo* after ligation. Then, transformation was subjected thereto by using *E. coli* XL1-Blue to
25 prepare cDNA library. The cDNA library gave clone pBDEXAM formed apparent halo due to decomposition of dextran in a medium containing blue dextran and IPTG.

It has been reported that *Lipomyces starkeyi* (hereinafter, abbreviated to as “*L. starkeyi*”) produces endodextranase (EC 3.2.1.11) that degrades dextran and α -amylase that degrades starch. The microorganism is applied in the field of food, but it has not been reported that the microorganism produces antibiotic material or other toxic material.

It has been reported that dextranases produced by microorganism are generally inducible enzymes except some bacterial dextranases. The present inventors have disclosed *L. starkeyi* ATCC 74054 which always produces both dextranase and amylase (U.S. Pat. 5,229,277) with characterizing the properties of the enzyme prepared by the microorganism that produces dextran of small size by using sucrose and starch. The present inventors also has a patent application pending that relates to a microorganism producing dexamase enzyme which can degrade dextran and starch at the same time by using *Lipomyces starkeyi* KFCC-11077, an enzyme of the microorganism, and a composition comprised of the enzyme.

The enzyme prepared from the gene (dexam) of the present invention degrades not only dextran and starch but also insoluble glucan. Hereinafter, the enzyme according to the present invention is referred to as “DEXAM enzyme”.

Dexamase of the present invention, in case that dextran is used as the substrate, essentially produces glucose, isomaltose and branched tetrose, and small amount of branched pentose and branched hexose. When starch is used as the substrate, the enzyme essentially produces glucose, maltose, maltotriose and maltotetrose, and various maltooligosaccharides as well.

Since DEXAM enzyme of the present invention can degrade levan (polymer of β -fructan), it can effectively degrade fructan, one of the component of plaque formation.

Thus, DEXAM enzyme of the present invention effectively degrades both

soluble glucan and insoluble glucan. In addition, the enzyme can repress formation of plaque or remove the plaque already formed by inhibiting aggregation of glucan and bacteria, thereby showing activity on preventing tooth decay.

From the experiments by using hydroxyapatite, which consists of similar components as tooth, DEXAM enzyme showed the property of retaining the bond to hydroxyapatite, being suggested that DEXAMase can be retained on tooth.

Further, the present invention relates to novel microorganism comprising gene encoding DEXAMase. The *E. coli* strain of the present invention was deposited in KCTC at Yuseong-gu, Daejeon on July 26, 2001 with Deposit No. KCTC 10024BP.

The present invention also relates to a process for preparing DEXAM enzyme. The process comprises the steps of culturing pBDEXAM clone and isolating DEXAM enzyme from the culture medium. As pBDEXAM can produce DEXAMase by using not only expensive dextran but also glucose, fructose, sucrose or starch, the process is economically advantageous. The properties of DEXAM enzyme obtained from pBDEXAM is substantially identical to those of DEXAMase isolated from *L. starkeyi* KFCC-11077.

In addition, the present invention relates to a composition for preventing tooth-decay that comprises DEXAM enzyme. The activity of the enzyme of the present invention can be retained in commercially available mouth washing solutions for a long time, with showing intense tolerance against enzyme inhibitors. Thus, the composition containing the enzyme of the present invention can be employed in various oral protective products such as toothpaste and mouth washing solution. The composition comprising the enzyme of the present invention can be applied to food such as chewing gum, soft drink or dairy products. The specific constituents of the composition may be appropriately determined by a person having ordinary skill in the technical field without difficulty.

Now, the present invention is described in more detail by referring to Examples which are not limitative:

In the present invention, the strain employed as DNA donor for preparing cDNA library was *Lipomyces starkeyi* KFCC 11077, which constitutively produces DEXAMase having dextranase and amylase activity. *Escherichia coli* XL1-Blue and λ -phage Uni-ZAP XR (Stratagene, USA) were employed as host cells and plasmid [phagemid] for preparing cDNA library.

L. starkeyi KFCC 11077 was cultured in LMS medium under aerobic condition at 28°C. The LMS medium comprised 1 %(w/v) of starch, 1%(v/v) of mineral solution and 0.3%(w/v) of yeast extract. Composition of the mineral solution was 2%(w/v) of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1%(w/v) of NaCl, 0.1%(w/v) of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1%(w/v) of $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, and 0.13%(w/v) of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$. As the medium for bacterial growth, LB (1% of tripton, 0.5% of yeast extract, 1% of NaCl, pH 7.3) and LBA (LB containing 50 μg of ampicillin/ml) were employed.

Example 1: Isolation of poly(A) + RNA from *L. starkeyi*

In LMS liquid medium, *L. starkeyi* KFCC-11077 was inoculated and cultured at 28°C for 36 hours. At the middle phase of exponential growth, the culture solution was centrifuged (6,500 xg) to obtain a cell pellet. The cell pellet was dissolved in a mixture of GIT solution [4M guanidine isothiocyanate, 25 mM sodium citrate (pH 7.0)], 0.5% lauroylsarcosyl dissolved in distilled water treated with 0.1% DEPC and 0.1 M 2-mercaptoethanol. Glass bead washed with acid and equal amount of phenol (pH 4.0) were added thereto and the mixture was subjected to vortex for 2 minutes. To the supernatant solution obtained after centrifugation, isopropanol was added to precipitate total RNA. From the oligotex-mRNA complex formed, mRNA was isolated by the use of oligotex resin. (Oligotex mRNA kit, Qigen).

Example 2: Preparation and screening of *L. starkeyi* cDNA

By the use of ZAP-cDNA Synthetic Kit (Stratagene, U.S.A.), cDNA was prepared from poly(A) + RNA (5 µg) obtained by culturing in a medium containing starch for 36 hours. The prepared cDNA was isolated as the size of not less than 500 bp by spin column separation. Then it was ligated with Uni-ZAP XR vector that had been treated with EcoRI-Xho. The ligated phage cDNA was subjected to packaging in vitro by using Gigapack Gold Kit (Stratagene, U.S.A.). The phage lysate thus obtained was transferred to pBluescript via excision in vivo, transformed with *E. coli* XL1-Blue, and cultured in LBA solid medium overnight. When recombinant colonies were grown up, top agar comprising 0.8% of agarose, 0.2% of blue dextran and 4 µl of 0.1M IPTG was placed on the plate, which was cultured at 28°C for 1-2 days to select the colonies in which blue dextran is hydrolysed to form obvious halo.

The selected colonies are named as pBDEXAM.

Example 3: Assay of base and amino acid sequence of cDNA clones

Plasmid DNA for assay of base sequence was prepared according to alkaline lysis method. The assay of base sequence was performed by means of GeneAmp 9600 thermal cylinder DNA sequencing system (model 373-18, Applied Biosystems, U.S.A.) and ABI PRISM Cycle Sequencing Kit (Perkin Elmer Corp. U.S.A.).

The base sequence of DNA fragment (5.4 kb) having glucanhydrolase gene had one open reading frame (ORF) comprised of 2679 bp of bases. Presumed amino acid sequence locates from the starting codon (ATG) at the 874th nucleotide position of the identified base sequence to the terminating codon (TGA) at the 3552nd position. The structural gene consists of 892 amino acids, and calculated molecular weight is 99.1 kDa.

Example 4: Selection of strain which expresses DEXAM gene

In order to confirm activity in the supernatant, clones which showed activity on the plate was subjected to isopropyl- α -D-thiogalactosidase (IPTG) induction. The selected colonies were cultured in 5 ml of LBA liquid medium at 37°C overnight. One percent (1%) (v/v) thereof was inoculated in LB medium again and cultured until it was 0.5 at OD_{600nm}, and IPTG was added to make the final concentration 0.1 mM. After culturing at 25°C overnight, the culture solution was centrifuged (6,500 xg, 15 minutes) to recover supernatant. After mixing the supernatant with 1% of dextranase or 1% of starch by 1:1, the mixture was reacted for 20 hours to confirm the activity.

Example 5: Preparation of coenzyme solution

Clone pBDEXAM was cultured in LBA liquid medium at 37°C overnight, of which 1% was inoculated again in LB medium to be cultured to make 0.5 at OD_{600nm}. IPTG was added until the final concentration became 0.1 mM, and the mixture was cultured at 25°C overnight. The culture solution was centrifuged (6500 xg, 15 minutes) and the supernatant solution was obtained to be used as the first enzyme solution. In order to concentrate the recovered first enzyme solution, enzyme not less than 30 kDa was concentrated by means of 30K cut-off hallow fiber (Milipore, U.S.A.). In order to remove the components of culture medium, it was concentrated by six times by exchanging four times with 20 mM citrate/phosphate buffer (pH 5.5) [2 L each time]. To the concentrate, polyethyleneglycol (PEG, MW=150,000 – 20,000) was added, and the mixture was reacted at 4 °C to concentrate by 20 times further. The PEG concentrate was dialyzed to 20 mM citrate/phosphate buffer (pH 5.5), and the dialyzed solution was used as the coenzyme solution for measuring protein activity.

Example 6 : Measurement of enzyme activity

In order to measure reducing value, DNS (3,5-dinitrosalicylic acid) method was employed. DNS reagent (100 μ l) and enzyme reaction mixture (50 μ l) and distilled water (50 μ l) were added, and the mixture was reacted at 80°C for 10 minutes, and chilled for about 15 minutes. Then, absorbance was measured at 550 nm. Dextranase activity of DEXAM enzyme was determined by measuring the amount of isomaltose produced after adding enzyme solution to buffer containing 2% dextran and reacting the mixture at 37°C for 15 minutes. One unit of dextranase is defined by the amount of enzyme that produces 1 μ M of isomaltose after reaction at 37°C for 1 minute by using dextran as substrate. Amylase activity is defined by the amount of enzyme that produces 1 μ M of maltose after adding the enzyme to buffer containing 2% of starch and reacting the mixture at 37 °C.

The enzyme activity of dextranase and amylase of the coenzyme solution (DEXAM) obtained from recombinant clone pBDEXAM was examined. As a result, it was confirmed that dextranase activity was 0.16 unit/ml and amylase activity was 0.25 unit/ml.

Example 7: Optimal activity and stability of DEXAM enzyme depending on pH or temperature

Optimal pH of dextranase and amylase activity of DEXAM enzyme was determined by reaction rate in the pH range of 4 – 9 (by interval of pH 1.0). For the experiment, 20 mM citrate-phosphate buffer (pH 4.0), citrate/phosphate buffer (pH 5-6) and sodium phosphate buffer (pH 7-9) were employed, and dextranase and amylase activity was determined by using DNS (3,5-dinitrosalicylic acid) method after reaction at 37°C for 48 hours. The pH stability of enzyme was measured after adding enzyme to each buffer and standing the mixture at 22°C for

48 hours.

Optimal temperature of enzyme activity was determined by measuring reaction rate after adding the enzyme at various temperatures (20-90°C, 10°C interval) and standing the mixture for 30 minutes. The temperature stability was obtained by measuring residual activity after placing the enzyme at various temperatures for 30 minutes.

Dextranase and amylase of DEXAMase showed optimal activity at pH 5.5. Amylase activity retained not less than 80% of optimal activity in the range of pH 5.5 – 7, while dextranase activity did in the range of pH 4.0 – 9.0 (Fig. 1, Table 2).

The temperature of maximum enzyme activity of dextranase and amylase was 40°C and 50°C, respectively. The dextranase activity and amylase activity retained 80% of initial activity at 50°C or lower temperature and 60°C or lower, respectively. (Fig. 2 and Table 3)

Example 8: Degrading properties of DEXAM enzyme on various substrate

Degradation of various substrate by the enzyme solution was examined. In addition to dextran, water-soluble starch and glucan, 1% (w/v) aqueous solutions of macromolecular polymers having various structures were prepared, and the mixture was mixed with enzyme concentrate in an equivalent amount to be reacted at 37°C for 5 days. The hydrolyzed product thus obtained was analyzed. For the conformation of hydrolysis of macromolecular polymers, levan (polysaccharide wherein D-fructose is bonded in the position of β -2,6), inulin (polysaccharide wherein D-fructose is bonded in the position of β -2,1) or mutan (polysaccharide wherein D-glucose is bonded in the position of α -1,3), or maltose (4-D- α -D-glucopyranosyl-D-glucose; disaccharide wherein two molecules of D-glucose are bonded in the position of α -1,4) or isomaltose (6-D- α -D-

glucopyranosyl-D-glucose; disaccharide wherein two molecules of D-glucose are bonded in the position of α -1,6) was employed.

Via reaction with dextran substrate, 14.0% of glucose, 31.2% of isomaltose and 33.5% of isomatotriose were essentially produced. Branched
5 oligosaccharides were produced as well. Via reaction with starch, glucose, maltose, isomaltotiose and maltotetrase were essentially produced in the amount of 28.2%, 18.4%, 24.7% and 28.6%, respectively. Thus, DEXAM is considered that it performs endo-dextranase and endo-amylase type reaction in the reaction with dextran and starch (Fig. 3).

10 In order to examine degradation property of DEXAM enzyme produced from clone pBDEXAM, macromolecular polymers having various structures are reacted with the enzyme. As a result, it showed hydrolytic properties on levan and inulin having β -bond as well as macromolecular polymers containing α -1,3-D-glucoside bond such as mutan. In addition, disaccharides such as maltose and
15 isomaltose were also hydrolyzed (Fig. 3).

Example 9: DEXAMase bond to hydroxyapatite (HA)

Having ability to directly attach to bones, calcium phosphate ceramics are widely used as alternative material of bones. Among them, hydroxyapatite (HA),
20 which has a lot of identical crystallographic properties to natural apatite existing in bones, is widely used as alternative material in research of artificial bones and teeth. Coherence of DEXAMase to HA was examined. Hydroxyapatite (Bio-Gel HTP, Bio-Rad Laboratories, Richmond CA) was suspended in 10 mM phosphate buffer (pH 6.8). DEXAMase was dissolved in the same buffer solution.
25 The enzyme activity was 10 U/ml. Each 200 μ l of HA and enzyme were mixed, and absorption was performed for 60 minutes. After washing unabsorbed enzyme, the adsorbed enzyme was gradually eluted by using 10, 50, 100, 200, 300,

400, 500 mM of phosphate buffer (pH 6.8) containing 1 M NaCl. Eluted enzyme fractions are collected, and dextranase and amylase activity was measured.

As can be shown in Fig. 4, dextranase and amylase of DEXAMase were eluted in 200 – 300 mM of hydroxyapatite. From the result, it is confirmed that DEXAMase has strong cohesion to hydroxyapatite, so that it remains on teeth for a long time.

Example 10: Effect of inhibitors on dextranase and amylase activity

Ethanol (5%) did not affect on activity of each enzyme, and sodium fluoride did not affect on amylase. However, EDTA reduced about 40% of initial activity of both enzymes, and 0.05% and 0.5% SDS reduced about 35~40% of the activity, while 10~20% ethanol reduced about 45% of dextranase activity and 15-30% of amylase activity (Table 4).

Example 11: Stability of dextranase and amylase against mouth washing components

Sodium benzoate gave no affect on dextranase and amylase activity of DEXAMase. In case of 0.01% of chlorhexidine, 90% of the activity was retained. However, in case of cetylpyridinium chloride, enzyme inactivity appeared depending on the concentration. When the concentration was 0.01%, about 90% or more activity was retained, while about 60% and 48% of dextranase and amylase activity was reduced, respectively, in case of 1% concentration (Table 5).

[Table 1] DEXAMase activity of pBDEXAM clone

Substrate	Before reaction	After reaction	OD ₅₅₀	Glucose (μg)	Enzyme activity (unit ^a /ml)
Dextran	0.046	0.089	0.043	13.21989	0.163209
Starch	0.042	0.205	0.163	20.11842	0.248376

In the Table, one unit^a of dextranase or alylase is defined as the amount of enzyme that can produce 1 μ mole of glucolse in an hour by using 2% (w/v) of dextran or water-soluble starch.

5

[Table 2] Effect of pH on DEXAMase activity and stability

	Dextranase	Amylase
Optimum pH	5.5	5.5
Stable pH range	4.0 – 9.0	5.5 – 7.0

In the Table, stable pH range means that residual activity is not less than 60% of the initial activity.

10

[Table 3] Effect of temperature on stability of DEXAMase

	Dextranase	Amylase
Stable temperature range (°C)	≤ 50	≤ 60

In the Table, stable temperature range means that residual activity is not less than 80% of the initial activity.

15 [Table 4] Dextranase activity and amylase activity of DEXAMase on various compounds

Inhibitor	Concentration	Relative activity (%)	
		Dextranase	Amylase
Control		100	100
EDTA	10 mM	68	60
SDS	0.05 %	62	75
	0.5 %	60	61
Fluorine	0.05 %	86	98
Ethanol	1 %	89	100
	5 %	100	100
	10 %	44	82
	20 %	46	71

[Table 5] Dextranase activity and amylase activity of pBDEXAM on chemical components used in mouth washing products

Chemical component	Concentration	Relative activity (%)	
		Dextranase	Amylase
Control		100	100
Sodium benzoate	1 %	99	100
Chlorhexidine	0.01 %	98	100
	0.02 %	71	100
	0.05 %	68	97
	0.1 %	83	94
Cetylpyridinium chloride	0.001 %	100	100
	0.01 %	98	92
	0.05 %	72	85
	0.1 %	54	71
	0.5 %	39	70
	1 %	40	52

5 As can be shown from the above constitution of the invention, DEXAMase produced by *Lipomyces starkeyi* KSM 22 according to the present invention is a single protein of about 100 kDa size, and shows both dextranase (EC 3.2.1.11) activity and amylase (EC 3.2.1.1) activity. As a result of assay of DNA base sequence, the base sequence of DNA segment (5.4 kb) had one open reading
10 frame (ORF) comprised of 2679 bp of bases. The gene of presumed structure consists of 892 amino acids, and the molecular weight was about 99.1 kDa.

As a result of confirming dextranase and amylase activity of coenzyme, dextranase activity was 0.16 unit/ml, while amylase activity was 0.25 unit/ml. Final reaction products of dextranase and amylase with dextran and starch were
15 typical products of endo-dextranase and endo-amylase. Dextranase activity essentially produced glucose (14.0%), isomaltose (31.2%) and branched tetrasaccharide (33.5%) as well as branched pentasaccharide. Amylase activity essentially produced glucose (28.2%), maltose (18.4%), matotriose (24.7%) and

maltotetraose (28.6%). In order to examine properties of DEXAMase to degrade various carbohydrates, the enzyme was reacted with macromolecular polymers of various structures. As a result, it showed the property of hydrolyzing macromolecular polymer comprising α -1,3-D-glucoside bond such as mutan as well as fructan comprising β -bond such as levan and inulin.


BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT
OF MICROORGANISMS FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued pursuant to Rule 7.1

TO : Lifenza Co., Ltd.
2nd Floor Samwoo Bldg.,
#732-24, Yeoksam-dong, Kangnam-gu, Seoul 135-514,
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: <i>Escherichia coli</i> XL1-Blue/pBDEXAM	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCTC 10024BP
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: [x] a scientific description [] a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on July 26 2001 .	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on _____ and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on _____	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Korean Collection for Type Cultures Address: Korea Research Institute of Bioscience and Biotechnology (KRIBB) #52, Oun-dong, Yusong-ku, Taejon 305-333, Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official(s):  BAE, Kyung Sook, Director Date: July 30 2001


WHAT IS CLAIMED IS:

1. An enzyme comprised of amino acid residue of sequence information 1, its mutant or a part thereof, which degrades insoluble glucan and has both dextranase activity and amylase activity.
2. Gene, mutant and/or segment of sequence information 2 which encodes the protein, its mutant or a part thereof claim 1.
3. The protein according to claim 1, wherein the mutant has not less than 80% of homology with the amino acid sequence of sequence information 1.
4. The DNA according to claim 1, wherein the mutant has not less than 80% of homology with the gene base sequence of sequence information 2.
5. The transformed cell which express DNA of claim 2.
6. The cell according to claim 5, which are prokaryotes or eukaryotes.
7. The cell according to claim 6, which are *E. coli* KCTC 10024BP.
8. A process for preparing enzyme degrading insoluble glucan and having both dextranase activity and amylase activity, which comprises the steps of (a) culturing cells of claim 5; (b) expressing enzyme from the cultured cells; and (c) purifying the enzyme.
9. Enzyme prepared by the process of claim 8.

10. A composition comprising the enzyme of claim 9.

1/5

FIG. 1a



2/5

FIG. 1b






FIG. 2



4/5

FIG. 3



5/5




FIG. 4

Sequence Listing

<110> KIM, DO MAN
LIFENZA CO., LTD.: KIM SUNG HYUK

<120> Enzyme with the removal activities of the plaques, DNA sequence encoding said enzyme, the expressing host cell and methods for producing and purifying said enzyme

<160> 2

<170> KopatentIn 1.71

<210> 1

<211> 892

<212> PRT

<213> E. coli XL1-Blue/pBDEXAM

<400> 1

Met Ala Thr Met Ala Asn Leu Leu Ala Leu Thr Leu Ser Ile Ser Lys
1 5 10 15

Phe Ala Ile Arg Ser Phe Met Asp Pro Thr Cys Asn Ser Cys Pro Gly
20 25 30

Cys Ser Ile Pro Val Pro Ile Trp Ile Pro Trp Ala Leu Gln Leu Gly
35 40 45

Ile Pro Ile Leu Pro Pro Ser Leu Tyr Leu Tyr Gly Ile Ser Val Gly
50 55 60

Ile Ser Ile Arg Lys His Leu Ser Asn Gln Arg Leu Pro Cys Ala Tyr
65 70 75 80

Leu Leu Gln Val Pro Cys Ser Ser Glu Pro Ser Cys Tyr Asn Asn Arg
85 90 95

His Gly Ser Cys Pro Leu Tyr Asp Ser Ile Leu Trp Ile Gly Ser Gly
100 105 110

Arg Asn Tyr Ala Asn Leu Ser Ser Gln Leu Val His Thr Trp Leu Val
115 120 125

Sequence Listing

Cys Trp Thr Val Asp Thr Gly Val Thr Pro Phe Tyr Ser Ala Leu Ile
 130 135 140

Gly Leu Thr Thr Gly Ser Ser Ile Tyr Arg Gln Ile Ser Gln Tyr Ala
 145 150 155 160

Thr Ile Ile Ile Gly Leu Gln Phe Leu Ala Gln Phe Leu Leu Asn Leu
 165 170 175

Leu Cys Asp Arg Cys Gly Val Val Gly Leu Ser Ser Phe Leu Gly Ser
 180 185 190

Ser Pro Met Arg Thr Ala Ala Met Gly Gln Leu Ser Ala Leu Ala Val
 195 200 205

Pro Leu His Pro Asn Gly Asn Ser Thr Gly Cys Cys Arg Ala Val Val
 210 215 220

Thr Leu Phe Gln Thr Arg Asn Ser Trp Pro Thr Arg Thr Thr Gly Leu
 225 230 235 240

Arg Lys Ser Leu Val Trp His Gly Lys Ser Arg Leu Tyr Thr Thr His
 245 250 255

Ala Tyr Thr Pro Phe Arg Ser Tyr Ser Tyr Arg Arg Leu Glu Cys Gln
 260 265 270

Val Ser Tyr Ser Thr Pro Thr Arg Cys Ile Pro Tyr Gly Ser Arg Ser
 275 280 285

Leu Met Gln Leu Asp Leu Glu Val Arg Ile His Ile Arg Leu Val Leu
 290 295 300

Ile Leu His Leu Pro Gly Leu Phe Ser Ser Leu Cys Val Gln Gly Ala
 305 310 315 320

Leu Ala Gly Val Thr Val Lys Gln Asn Asn Ala Ala Ala His Ala Ile
 325 330 335

Sequence Listing

Leu Ser Leu His Gly Tyr Val Val Lys Val Asn Ala Glu Asp Asn Trp
 340 345 350

Ile Ala Val Ile Cys Asp Ser Pro Asn Gln Pro Met Trp Tyr Thr Ser
 355 360 365

Ile Leu Thr Gly Ile Gln Ala Trp Asn Cys Gly Gly Lys Thr Ile Asn
 370 375 380

Ala Gln Pro Ser Lys Thr Ile Tyr Phe Asp Ala Thr Thr Gly Leu Val
 385 390 395 400

Trp Val Ala Ser Arg Val Ser Thr Arg Trp Lys Ile Trp Met His Asn
 405 410 415

Leu Arg Ile Leu Thr His Arg Gln Arg Thr Leu His Pro Tyr Glu Thr
 420 425 430

Gly Thr Leu Tyr Ile Ser Gly Ile Gly Asp Phe Tyr Asp Leu Thr Glu
 435 440 445

Thr Ser His Cys Gln Glu Ser Met Thr Pro His Ile Ala Asp Leu Asp
 450 455 460

Pro Ala Leu Pro Gly Ser Lys Phe Cys Ala Pro Trp Pro Gln Ser Gln
 465 470 475 480

Lys Pro Asn Leu Trp Ile Arg Gly Gly Thr Tyr Leu His His Phe Val
 485 490 495

Pro Ala Gly Thr Leu Ala Leu Gly His His Val Ser Val Gly Ser Thr
 500 505 510

Ser Ser Asp Ala Ala Ile Phe Ser Arg Ile Val Ser Cys Trp Arg Val
 515 520 525

Val Leu Ala Val Tyr Cys Val Ser Phe Ile Ala Ala Arg Ser Arg Asn
 530 535 540

Ile Val Glu Tyr Cys Ala Ala Arg Gln Phe Pro Ser Met Met Met Gln

Sequence Listing

545	550	555	560
Asn Pro Ser Trp Cys Thr Trp Glu Gln Asp Ala Pro Pro Trp Pro Ala			
	565	570	575
Ala Thr Cys Tyr Ser Ile Thr Ser Ile Ile Ala Pro Asp Gly Ile Tyr			
	580	585	590
Ile Pro Gly Lys Ile Ser Gly Val Val Met His Glu Pro Ile Gly Val			
	595	600	605
Ile His Thr Lys Tyr Ile Lys Asp Gly Thr Ala Phe Gly Tyr His Trp			
	610	615	620
Pro Leu Trp Ile Leu Cys Asn Leu Asp Pro Ile Tyr Asn Cys Ser Ser			
	625	630	635 640
Arg Tyr Arg Asn Val Val Cys Glu Leu Leu Cys Pro Thr Leu Phe Arg			
	645	650	655
Cys Ser Ser His Pro Ala Ala Ile Asp Asn Arg His Asp Val Cys Gly			
	660	665	670
Arg Leu Leu Ala Ser Ser Arg His Arg Phe Leu Ser Arg Val Leu Lys			
	675	680	685
Phe Gly Thr Ala Glu Ser Leu Thr Arg Tyr Gln Val Glu Asn Phe Ile			
	690	695	700
Val Lys Arg Gln Val Asp Ile Leu Arg Val Asp Leu Leu Arg Pro Arg			
	705	710	715 720
Pro Ser Asp Gln Arg Leu Ser Glu Ser Ala Met Ala Ala Arg Val Gly			
	725	730	735
Ala Ala Lys Tyr Arg Arg Arg Leu Ile Arg Ala Lys Ser Arg Leu Tyr			
	740	745	750
Ala Thr Gln Lys Met Ile Glu Asp Arg Glu Met Glu Ile Ala Arg Leu			
	755	760	765

Sequence Listing

Lys Gln Arg Ile Arg Glu Ser Arg Leu His Arg Asp Ser Thr Arg Ser
 770 775 780
 Gly Phe Ala Arg Tyr Gly Gln Asn His Gly Ser Lys His Arg Ala Gly
 785 790 795 800
 Asn Asn Asp Gly Glu Gly Ala Leu Leu Leu Phe Lys Asn Val Val Val
 805 810 815
 Pro Asp Thr Gly Tyr Thr Asn Ile Cys Trp His Arg Arg Asn Ser Leu
 820 825 830
 Ser Ser Thr His His Cys Leu Thr Thr Arg Thr Gly Arg Ile Ser Pro
 835 840 845
 Thr Leu Trp Leu Val Ala Ala Thr Lys Ser Val Tyr Asn Phe Lys Pro
 850 855 860
 Cys Arg Gln Ser Pro Gly Ala Tyr Trp Arg Val Asp Ala Ala Ile Leu
 865 870 875 880
 Leu Ile Ser Ile Ala Trp Gly Glu Pro Glu Ser Thr
 885 890

<210> 2
 <211> 5422
 <212> DNA
 <213> E. coli XL1-Blue/pBDEXAM

<400> 2
 cgcgcttcca ggcctgttg ctgggtatcg tcttcagca gatcgcgag gttaacgccg 60
 ccttcaccaa acaggcagag atggagtta ccaatggagg aaacgcgcag gcggttgcaa 120
 gtggcgcaga agtctttttc atacggcatg ataaggccaa tctctccggc gtaatctgga 180
 tggcaaaaga cttgcgcggg accgtcgctg cggttgacgta attggtggat ccagccgcga 240

Sequence Listing

cgtagtagct cgtcacgcag aacctgacca gagatgtgat gcttacggaa gagctcgtg	300
ccctcgcccg ttccatcag ttcatgaaa cgcagctgga tagggcgatg ctggatccag	360
ttcagaaagg tgtagagctg gtgatgatta acatcacgca tcagcacggt attgactttg	420
accttctcaa aaccggcctc aaatgcagca tcaatccctg ccatgacctg gttgaattta	480
tcctgcccgg taatagcgtg aaactggcgg gcgtccagac tgcgacact gacgttaatg	540
ccagtaagtc ccgcatcgcg ccagctcgcc acatcgcggt ccagacggta accattgggt	600
gtgaccgcaa tctggcaatt agcgtcgttt tccgcacag cggcgataac gaccaacaat	660
gggatgctgt .gaaacaaatt gtgctaaaac gatgcaagct attcaatata tgaagtgact	720
acttatagct ttgcctaaaa tgaaggctct agtatatgcg ttagcacaat atggtttgag	780
tcctcatcta cctataagaa tatctcttag tctttcgggt gaaatccttc ggactgtctc	840
tctttgcgtt gaagaagggt taatctcaag cctatggcca caatggcaaa cctacttgcc	900
ttgacgttt caattagcaa gtttgctatt cgatcattca tggaccaaac ttgcaattct	960
tgtcccggtt gttccatccc ggtaccata tggataccat gggcactaca attgggaata	1020
cccattttgc cgccgagctt gtacctctat ggcactctcg tgggaatatc gatacgcaaa	1080
cacctgtcca accagcgatt gccgtgcgca tatctcttac aggtaccgtg cagttcggaa	1140
ccaagctgct acaacaatcg tcatggctcc tgccccctat atgattcgat cctatggatc	1200
ggaagtggtc gcaactacgc caacctatcc agccaactgg tgcatacgtg gctcgtgtgc	1260
tggactgtcg aactggcgt taccctgttt tactcggcct taataggcct tactactggg	1320
tcgagtattt accgacaaat ctctcagtae gccacgatca tcattggact gcaattcttg	1380
gcccagtttc ttctcaatct cctatgcgac aggtgtggcg tgggtggatt gtcaccttt	1440

Sequence Listing

cttggatcct cgccgatgcg aacggcggca atgggtcaat tgagtgcatt ggctgtccca	1500
ttacacccca atggaaattc tactggatgt tgtcgagcag tggtcacatt atttcagact	1560
cgaaattctt ggcctacacg cactactggt cticgcaagt ccttagtctg gcatggaaag	1620
agtcgtcttt atacgacaca cgcttataca ccattcaggt cctactctta tcggcgactg	1680
gagtgccaaag tcagctattc tactcccacc aggtgtatac cctatggatc aagatcattg	1740
atgcaactcg atttagaagt tcgtattcat atacgtctag ttttaatttt acacttaccc	1800
ggcttgttct cgagcctgtg cgtacaaggt gcccttgctg gagtaactgt caagcagaac	1860
aatgcagctg cacatgcgat cttatcgctt catggctacg ttgtcaaagt gaatgccgaa	1920
gacaactgga tagcagtcac ttgcgattct ccgaaccagc cgatgtggta taccagtatc	1980
cttaccggta ttcaagcctg gaactgcggt ggaaagacca tcaatgcgca accaagcaag	2040
actatatatt tcgatgctac tacaggtctt gtctgggtgg cttcacgagt cagcaccagg	2100
tggaaaattt ggatgcacaa tcttcgtatc ctgaccacaa gacagcgaac ttgacacccc	2160
tacgaaacag gtacgtctta tatcagtggc attggcgatt tctacgactt gaccgaaacc	2220
agtcactgcc aagagtcaat gactccacat attgctgatt tagaccctgc tcttcggga	2280
tccaaatttt gcgccccttg gcccagagc cagaagccca atttgtggat ccgaggtggc	2340
acttacctcc accacttcgt ccctgcgggc acccttgccc tcgggcatca cgtctcagt	2400
ggatctacaa gctcggacgc tgctatattc agtcgtattg tgcctgctg gagagtcgtc	2460
ttagctgtct attgcgtgag cttcatagct gctcgctcca gaaatatcgt agaatactgt	2520
gctgcacgac aatttcgctc aatgatgatg caaaatcctt cctgggtgtac gtgggagcag	2580
gatgtccgc cgtggcctgc tgctacctgc tattccatta cgtcaatcat agctccagat	2640

Sequence Listing

ggtatttaca ttccgggcaa aatcagtga gtagtaatgc acgaaccaat aggtgttatt	2700
cacaccaa atcatcaaaga cggctactgcc ttcggctacc attggcccct ctggattcta	2760
tgcaacttgg atccaatcta caactgtcta tcacggtaic gaaacgttgt ttgcgaactg	2820
ctttgcccga ctctattccg ctgctcatca caacctgtctg ctattgacaa tcgacacgac	2880
gtctgcggtc gtctcttagc ttcttcacgg caccggtttt tgtccagggt actgaagttt	2940
ggaactgcag aatcacttac acggtatcag gtcgagaact tcatagtaaa acggcaagta	3000
gatataattac gtgtcgatct attgcggcca cgaccgtcag atcagagact gtcagaaagt	3060
gcaatggcag cgagggtcgg agctgccaaag tatcggcggc ggctcattcg ggccaagagt	3120
cgtctttatg cgacacaaaa gatgatcgag gacagagaaa tggagattgc tcgcctcaaa	3180
cagcgaatac gagagagccg actacatcga gattcaactc gttctggttt tgctcgttat	3240
ggtcaaaatc atggcagcaa acatcgcgct gggaacaacg acggcgaagg tgctttgttg	3300
ctattcaaaa atgtcgtcgt cccagacact ggctacacga atatctgctg gcacaggaga	3360
aactcattat ccagcacgca tcaactgtcta acgactcgga ctggacgtat atctccaact	3420
ctgtggttgg tggctgctac aaaaagcgtg tacaacttca agccgtgtcg tcaatctcct	3480
ggggcatatt ggcgcgtcga tgcagctatc ctgctgatat ccattgcgtg gggggagcca	3540
gagtctacgt gaattcagcc tccctgctcg cgtgagtgcg tgtgaccctt tctttaccac	3600
gctggagaga taatcggcta agacgttttag catgaaggcg attctcgctc gattctctct	3660
acataggcat attagttagc tcatgtgaag gcgatttcta gttcattctc taagctggaa	3720
cttattacct gttcgggcat tgtgacactc aacttgcctt caataagacg tggtcgatag	3780
gttacggtag atcagtctga ggtacgtcaa caatcactta ataagcacga gttggcaggt	3840

Sequence Listing

ttccttactc tgtcttgcac gcctaactcc aatttcgtgc attacaggat ccgccgtgta	3900
cttccaggga aaatctgaaa tgggactcag ggctaaagcg agaaccttaa gtcaatagat	3960
attatgcac tgcaggggct ggtatgatct ccagctatca gggcactaat acatttgcaa	4020
gtacgaaag acgagataga cgacgaatgg acgaataatc acgcattact ccacaggctt	4080
ccacggaggc tcagggiccc aacgatatct aggtcagacg gtatgatacc gcaccgatct	4140
gtattgcagt aactatctcc gtcagctac aggcagcgca cgtataatgg tggatggcat	4200
gcagagaatt ccagccgtag ctgtttcaca atgcaatatt acgcaagtga aattaacat	4260
cattaaagga ccaaagtga gtccaccgcg ttgcggccgc tctagaacta gtggatcccc	4320
cgcagcgccg ttgaaatacg cgcttccaag cgcctgttgc tgggatcgtc ttccagcaga	4380
tcgcgaggt taacgccgccc ttaccaaac aggcagagat ggagtttacc aatggaggaa	4440
gcttacgacg cgcaggcggc tgcaagtggc gcagaacctg accagagatg tgatgaagag	4500
ctcgctgccc tcgcccgttt ccatcagttc gatgaaacgc agctggatac cctgccatga	4560
cctggttgaa tttatcctgc ccggtaatag cgtgaaactg gcgggcgtcc agactgtcga	4620
cactgacgtt aatgccagta agtcccgcac cgcgccagct cgccacatcg cgttccagac	4680
ggtaaccatt gggtgtgacc gcaatctggc ggatagcgtc gttttccgc acagcggcga	4740
tgatatcggc aaagtcgcgg cgtaaagacg gctctcctcc tgtcaggcgc actttttcgg	4800
gtgcccagtc tggcgaaggc gcgcgtaacc cggcgaattt catcgacggt aagaaagcct	4860
ttattggatga cgccgctcgg ttgttagcca tccggcagcc agtaggtgca acgaaagtta	4920
cacacatcgg taatcgacag gcgcaagtag taaaacttac gcgcaaatgc atcagtcagt	4980
tgtgaagcca tgtacacctt tccagatacg ggaggcgaag tcattttctt cttcgccctg	5040

Sequence Listing

gtggcaatct ttacgagcat tgtcacggcc aaagcacctg atcagttgac ccaggtgcag	5100
aggctagagt gtttagtggt tatgccgata ctagcgtgta aatgagattt ttaccatcca	5160
cactttcgct atgtaatcat gtatatagcg tcatgatcgt gctatttcgc tacagaataa	5220
gggaaaatgg cggaagaaat cgaagcgttg atggttgctg cgagaaagca atgagcgtaa	5280
tcatcgtcgg tggcggcatg gcgggcgcga cgctggcgct ggctatttcc cggtaaagtc	5340
acggggcgct gccggtacat ttgattgaag cgactgcgcc agagtcacat gctcatccgg	5400
gctttgatgg acgagcgata ga	5422

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR01/01439

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C12N 9/24, C12N 15/56, C12N 1/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N 9/24, C12N 15/56, C12N 1/20

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
GenBank, PubMed, CA, Espacenet, PAJ, USPTO, "Lipomyces starkeyi", "amylase", "dextranase"

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Kim, D. et al., "Characterization of a novel carbohydrase from <i>Lipomyces starkeyi</i> KSM 22 for dental application", J. Microbiol. Biotechnol., 9(3), 260-64, 1999.	1-10
X	Ryu, S.J. et al., "Purification and partial characterization of a novel glucanhydrolase from <i>Lipomyces starkeyi</i> KSM 22 and its use for inhibition of insoluble glucan formation", Biosci. Biotechnol. Biochem., 64(2), 223-8, 2000.	1-10
E	WO 01/66570 A1 (Kim, D. et al.), 13 Sep. 2001.	1-10
A	Kelly, C.T. et al., "Thermostable extracellular alpha-amylase and alpha-glucosidase of <i>Lipomyces starkeyi</i> ", Appl. Microbiol. Biotechnol., 22(5), 352-8, 1985.	1-10
A	Koenig, D.W. et al., "Induction of <i>Lipomyces starkeyi</i> dextranase", Appl. Environ. Microbiol., 55(8), 2079-81, 1989.	1-10

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 MAY 2002 (21.05.2002)

Date of mailing of the international search report

23 MAY 2002 (23.05.2002)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
Government Complex-Daejeon, 920 Dunsan-dong, Seo-gu,
Daejeon Metropolitan City 302-701, Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

LEE, Cheo Young

Telephone No. 82-42-481-5594



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.